



**General Certificate of Education**

**Chemistry 6421**

**CHM5      Thermodynamics and Further  
Inorganic Chemistry**

**Mark Scheme**

*2008 examination - June series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: [www.aqa.org.uk](http://www.aqa.org.uk)

Copyright © 2008 AQA and its licensors. All rights reserved.

#### COPYRIGHT


AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.


---

**CHM5****SECTION A****Question 1**

- (a) Proton ( or  $H^+$ ) acceptor (1)
- (b) Electron (or lone) pair donor (1)
- (c) Electron (or lone) pair donor (1)  
(Ignore answers that talk about attraction to +ve centre)  
(allow Lewis base)
- (d)  $NH_3 + H^+ \rightarrow NH_4^+$  (1)  
(or  $NH_3 + H_3O^+ \rightarrow NH_4^+ + H_2O$ )  
(allow  $Cl^-$  as a spectator)
- (e)  $4NH_3 + [Cu(H_2O)_6]^{2+} \rightarrow [Cu(NH_3)_4(H_2O)_2]^{2+} + 4H_2O$   
Correct copper species (both) (1)  
(allow no square brackets or other shapes of brackets)  
balanced equation (1)  
(only with correct species)  
colour of reagent: Blue (1)  
Colour of product: (Dark) blue (1)  
(note NOT purple, NOT blue ppt)  
(Note mark colours independently correct)



(nucleophilic) addition-elimination (1)



(final  $\text{Cl}^-$  not essential)

(ignore final proton donation to base even if arrow etc wrong)

arrow from lone pair on ammonia to C (1)

arrow from  $\text{C}=\text{O}$  to O (1)

intermediate with + and – charges (1)

3 arrows and lone pair on O (1)

**Total 14 marks**

**Question 2**

(a)  $\Delta G = \Delta H - T\Delta S$  (1)

(b) (Boiling is a) spontaneous (or feasible) (change) (1)  
(or (water and water vapour are at) equilibrium)

(c) When  $\Delta G = 0$   $\Delta S = \Delta H/T$  (1)

$$= 23.4 \times 1000/240$$

$$= 97.5 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)} \quad (1)$$

(units not essential but 97.5 with wrong units scores 1/2)

(note 0.0975 (kJ K<sup>-1</sup> mol<sup>-1</sup>) scores 1/2)

(allow 2 marks for correct answer)

(allow range 97 to 98)

(note, if -ve answer, can score first mark only)

(d) H bonding in both (1)

Stronger in HF (1)

(or more energy needed to overcome forces)

Because H—F is more polar than H—N (1)

(or electronegativity of F > N)

(or F is more electronegative or F is the most electronegative)

Note

(if breaking covalent bonds or ionic bonds C.E. = 0/3)

(allow 1/3 (second mark) for intermolecular forces in HF stronger without specifying nature of force or when comparing H bonding in HF with dipole-dipole or VdW in NH<sub>3</sub>)

**Total 7 marks**

**Question 3**

(must be this equation not a multiple)

(ignore state symbols)

(b)  $\Delta S = \Sigma S(\text{products}) - \Sigma S(\text{reactants})$  (1)

(must have  $\Sigma$  (or equivalent) and no  $\Delta$  on RHS)

$= 193 - (192/2 + 3/2 \times 131)$  (1)

(this also scores first mark)

$= -99.5 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$  (1)

(units not essential but penalise wrong units one mark)

(allow 3 for correct answer)

(allow range -99 to -100)

(if equation doubled allow 2/3 for -198 to -200)

(allow 1/3 for +99.5)

(can only score 1/3 (first mark) if answer is -130 and equation stated correctly)

- (c) (i)  $\Delta G = \Delta H - T\Delta S$   
 $= -46.2 - (700 \times -99.5)/1000$  (1)  
(or  $= -46.2 - (700 \times x)/1000$  if using given value or value from (b))  
 $= + 23.45 \text{ kJ mol}^{-1}$  (1)  
(allow range 23 to 24)  
(units must be given, penalise wrong units)
- Allow 2 for consequential marking from answer to (b) e.g.  
(if answer to (b) is +99.5 allow -115 to -116)  
(if answer to (b) is -199 allow 46 to 47 or 93 to 94)  
(if answer to (b) is -130 allow 44 to 45)  
(if used given answer of -125 allow 41 to 42)
- (ii) Decreases (or becomes more negative) (1)
- (d) To speed up reaction (1)  
(or fast reaction)  
(or give more molecules  $E > E_a$ )

**Total 8 marks**

**Question 4**

- (a) (i) W Pt (or in words) (1)
- X KCl, NH<sub>4</sub>Cl etc (allow any simple soluble salt and ignore water, paper, agar etc) (1)
- Y Mg (1)
- Z MgCl<sub>2</sub> (1)
- (aq not essential)
- (allow any identified soluble Mg salt)
- (ii) Pt|H<sub>2</sub>(g)|H<sup>+</sup>(aq)||Mg<sup>2+</sup>(aq)|Mg (1)
- (allow Mg|Mg<sup>2+</sup>(aq)||H<sup>+</sup>(aq)|H<sub>2</sub>|Pt ) (1)
- Species
- (ignore state symbols)
- (allow any coefficients)
- Correct order (1)
- (order is consequential on correct species)
- (can score this mark (not first mark) if phase boundary solidus omitted)
- (If Pt omitted max 1)
- (b) (i) 0.84 (V) (1)
- (ii) (+)3 (1)
- (or III)
- (or Mn<sup>3+</sup> or Mn(III))
- (iii) 2MnO<sub>2</sub> + 2H<sub>2</sub>O + Zn → 2MnO(OH) + 2OH<sup>-</sup> + Zn<sup>2+</sup> (1)
- (allow multiples)
- (allow Zn(OH)<sub>2</sub>)
- (arrow can be equilibrium arrow)
- (iv) *Oxidising agent* MnO<sub>2</sub> (1)
- (allow in words manganese oxide)
- Reducing agent* Zn (1)

- (v) Zn (or MnO<sub>2</sub>) used up (1)  
(or concentration of products increases)  
(or electrode(s) worn away)  
(allow polarisation or explanation in terms of ion migration)  
(note if equation reversed allow conseq i.e. Zn<sup>2+</sup> or MnO(OH) used up)



(or  $2\text{H}^+ + \text{H}_2\text{SO}_4$  etc)



(or  $2\text{H}_2\text{SO}_4 + 2\text{KBr} \rightarrow \text{K}_2\text{SO}_4 + \text{SO}_2 + 2\text{H}_2\text{O} + \text{Br}_2$ )

(allow production of SO<sub>3</sub><sup>2-</sup> for last mark but not for half equation i.e. 1/2)

- (ii) H<sub>2</sub>SO<sub>4</sub> cannot oxidise Cl<sup>-</sup> (1)

(or Cl<sup>-</sup> ions (or KCl) cannot reduce H<sub>2</sub>SO<sub>4</sub>)

(or Cl<sub>2</sub> strong(er) oxidising agent (than H<sub>2</sub>SO<sub>4</sub>))

(or Cl<sup>-</sup> weak reducing agent)

(allow any correct E<sup>0</sup> argument)



(or  $\text{H}_2\text{SO}_4 + 2\text{KCl} \rightarrow \text{K}_2\text{SO}_4 + 2\text{HCl}$ )

(or  $\text{H}^+ + \text{Cl}^- \rightarrow \text{HCl}$  or any correct equation to give HCl)

**Total 17 marks**

**Question 5**

- (a) Curve Y starts at origin and is steeper than curve A (1)  
Finishes at the same level as curve A (1)
- (b) Curve X starts at the origin and is below curve B (1)  
Approaches the same level as curve B (1)
- (c) Order is 1 (or first order) (1)  
(Note C.E. if order not equal to 1)  
When concentration (of iodine) is doubled gradient (or rate) doubles (1)  
(or when concentration (of iodine) is halved gradient (or rate) halves
- (d) (i)  $\text{S}_2\text{O}_8^{2-} + 2\text{Fe}^{2+} \rightarrow 2\text{SO}_4^{2-} + 2\text{Fe}^{3+}$  (1)  
 $2\text{Fe}^{3+} + 2\text{I}^- \rightarrow 2\text{Fe}^{2+} + \text{I}_2$  (either order) (1)  
  
(allow correct equations that are not ionic)
- (ii) Alternative route (1)  
Not used up (or is regenerated) (or not chemically changed) (or not in overall equation) (1)  
Speeds up reaction (or changes rate)  
Lowers activation energy  
(any two of these four)

- 
- (e) (i) Different phase (or state) from reactants (1)  
(or implied eg silver is a solid, reactants are gases)
- (ii) Reactants adsorb weakly (or poorly) (onto surface of silver) QWC mark (1)
- (iii) Reaction may be too fast (1)  
(note candidates must give the idea of reaction rate)
- Explosion (1)  
(or uncontrolled)  
(note do not accept further oxidation arguments)

**Total 14 marks**

## SECTION B

## Question 6

- (a) (i) Note incorrect reagent (e.g.  $\text{BaCO}_3$ ) CE = 0 but if  $\text{Ba}^{2+}$  or  $\text{Ba}^+$  implied, lose reagent mark and mark on

If two reagents given (one for each member of pair), mark first and ignore second

<b>Reagent</b>	$\text{BaCl}_2/\text{H}^+$ or $\text{Ba}(\text{NO}_3)_2$	$\text{Ba}(\text{OH})_2$	Ba	(1)
----------------	-------------------------------------------------------------	--------------------------	----	-----

<i>Obs with <math>\text{CuSO}_4</math></i>	(White) ppt	White and blue ppts	White and blue ppts	(1)
--------------------------------------------	-------------	------------------------	------------------------	-----

<i>Obs with <math>\text{Cu}(\text{NO}_3)_2</math></i>	No change or green or yellow solution	Blue ppt	Blue ppt	(1)
-------------------------------------------------------	---------------------------------------------	----------	----------	-----

$\text{CuSO}_4(\text{aq}) + \text{BaCl}_2 \rightarrow \text{BaSO}_4(\text{s}) + \text{CuCl}_2(\text{aq})$  (1)

(or  $\text{Ba}^{2+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{BaSO}_4(\text{s})$ )

(ignore state symbols)

(If use Ba, also need an equation to show production of  $\text{Ba}(\text{OH})_2$  or  $\text{Ba}^{2+}$ )

- (ii) If reagent incompletely given (e.g.  $\text{OH}^-$ ), lose reagent mark and mark on

<b>Reagent</b>	NaOH	xs NaOH	$\text{NH}_3$	xs $\text{NH}_3$	$\text{Na}_2\text{CO}_3$	(1)
----------------	------	---------	---------------	------------------	--------------------------	-----

(or conc) (or  
 $\text{NaHCO}_3$ )

<i>Obs with <math>\text{CrCl}_3</math></i>	Green ppt	Green solution	Green ppt	Purple solution	Green ppt gas evolved	(1)
--------------------------------------------	-----------	-------------------	-----------	--------------------	--------------------------	-----

<i>Obs with <math>\text{FeCl}_2</math></i>	Green ppt goes brown on standing	Green ppt	Green ppt goes brown on standing	Green ppt	Green ppt or white ppt	(1)
--------------------------------------------	----------------------------------------	-----------	----------------------------------------	-----------	---------------------------	-----

Note other answers possible e.g.  $\text{Zn}/\text{HCl}$  (1) blue solution (1) no reaction (1)

*Equations for reactions with  $\text{CrCl}_3$*  (Note square brackets for complexes & ss optional) (1)

NaOH  $\text{Cr}(\text{H}_2\text{O})_6^{3+} + 3\text{OH}^- \rightarrow \text{Cr}(\text{H}_2\text{O})_3(\text{OH})_3 + 3\text{H}_2\text{O}$

(or  $\text{CrCl}_3 + 3\text{OH}^- \rightarrow \text{Cr}(\text{OH})_3 + 3\text{Cl}^-$ ) etc

xs  
NaOH  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+} + 6\text{OH}^- \rightarrow [\text{Cr}(\text{OH})_6]^{3-} + 6\text{H}_2\text{O}$

(or  $\text{CrCl}_3 + 6\text{NaOH} \rightarrow \text{Cr}(\text{OH})_6^{3-} + 6\text{Na}^+ + 3\text{Cl}^-$ )

(allow formation of  $[\text{Cr}(\text{H}_2\text{O})_2(\text{OH})_4]^-$  &  $[\text{Cr}(\text{H}_2\text{O})(\text{OH})_5]^{2-}$ ,

NH<sub>3</sub> As NaOH but can have + NH<sub>3</sub> → NH<sub>4</sub><sup>+</sup> instead of + OH<sup>-</sup> → H<sub>2</sub>O

xs NH<sub>3</sub> [Cr(H<sub>2</sub>O)<sub>6</sub>]<sup>3+</sup> + 6NH<sub>3</sub> → [Cr(NH<sub>3</sub>)<sub>6</sub>]<sup>3+</sup> + 6H<sub>2</sub>O

Na<sub>2</sub>CO<sub>3</sub> 2[Cr(H<sub>2</sub>O)<sub>6</sub>]<sup>3+</sup> + 3CO<sub>3</sub><sup>2-</sup> → 2Cr(H<sub>2</sub>O)<sub>3</sub>(OH)<sub>3</sub> + 3CO<sub>2</sub> + 3H<sub>2</sub>O

NaHCO<sub>3</sub> [Cr(H<sub>2</sub>O)<sub>6</sub>]<sup>3+</sup> + 3HCO<sub>3</sub><sup>2-</sup> → Cr(H<sub>2</sub>O)<sub>3</sub>(OH)<sub>3</sub> + 3CO<sub>2</sub> + 3H<sub>2</sub>O

*Equations for reactions with FeCl<sub>2</sub>* (1)

NaOH [Fe(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup> + 2OH<sup>-</sup> → Fe(H<sub>2</sub>O)<sub>4</sub>(OH)<sub>2</sub> + 2H<sub>2</sub>O

(& xs) (or FeCl<sub>2</sub> + 2NaOH → Fe(OH)<sub>2</sub> + 2NaCl)

NH<sub>3</sub> & xs As NaOH but can have + NH<sub>3</sub> → NH<sub>4</sub><sup>+</sup> instead of + OH<sup>-</sup> → H<sub>2</sub>O

Na<sub>2</sub>CO<sub>3</sub> Fe<sup>2+</sup> + CO<sub>3</sub><sup>2-</sup> → FeCO<sub>3</sub>

(or FeCl<sub>2</sub> + Na<sub>2</sub>CO<sub>3</sub> → FeCO<sub>3</sub> + 2NaCl)

NaHCO<sub>3</sub> As NaOH or Na<sub>2</sub>CO<sub>3</sub>

(b) (i) 2MnO<sub>4</sub><sup>-</sup> + 16H<sup>+</sup> + 5C<sub>2</sub>O<sub>4</sub><sup>2-</sup> → 10CO<sub>2</sub> + 8H<sub>2</sub>O + 2Mn<sup>2+</sup> (1)

(ii) Moles C<sub>2</sub>O<sub>4</sub><sup>2-</sup> = vol in dm<sup>3</sup> × conc = 17.6/1000 × 0.1 = 0.00176 (this answer only) (1)

Moles MnO<sub>4</sub><sup>-</sup> = 2/5 × moles C<sub>2</sub>O<sub>4</sub><sup>2-</sup> (this mark is for 2/5) (1)

= 2/5 × 0.00176 = 0.000704 (or 7.04 × 10<sup>-4</sup>) (1)

(This answer only which also scores the previous 2 marks)

(iii) Mass of 1 mol of unknown = 0.1/0.000704 = 142 (1)

(or if M<sub>r</sub> assumed, mass of 1.0 g (or 0.1 g for 25 cm<sup>3</sup>) can be calculated from no. of moles × M<sub>r</sub>)


(must show working using answer from (b) (ii) to score this mark)

Unknown corresponds to NaMnO<sub>4</sub> (1)


(this mark only given if previous mark for working also given)

**Total 15 marks**


## Question 7



Trigonal planar (allow triangular planar) (1)




tetrahedral (1)



Square planar (allow octahedral if lone pairs shown) (1)




linear (1)




(note if more than one ligand shown, all must be correct)

(Second mark only given if first mark gained)



(Equal chance of) attack on each side of carbon (or molecule or double bond) (1)

(allow from above and below the plane)



Note, do not allow structures with bond angle of 90°

Allow CN or NC linkages as above

**Total 15 marks**

**Question 8**

(a)  $\Delta H_{\text{sol}}^{\ominus} = \Delta H_{\text{lattice}} + \sum \Delta H_{\text{hyd}}^{\ominus}$  so  $\Delta H_{\text{lattice}} = \Delta H_{\text{sol}}^{\ominus} - \sum \Delta H_{\text{hyd}}^{\ominus}$  (or cycle) (1)

For NaCl = +3.9 + 406 + 364 = (+) 774 (kJ mol<sup>-1</sup>) (1)

(allow 773.5 to 774)

For MgCl<sub>2</sub> = -155 + 1920 + 728 = (+) 2493 (kJ mol<sup>-1</sup>) (1)

If either of last two answers is correct first mark is also scored

(if both answers numerically correct but negative signs allow 1/3)

(b) Attraction (or force or bonding) between ions weaker (ions for QWC) (1)

(or ionic bonding weaker)

Charge on Na<sup>(+)</sup> or less than that on Mg<sup>(2+)</sup> (1)

(Do not allow polarisation argument)

(c) Al<sup>(3+)</sup> ions) have higher charge/size ratio than Mg<sup>(2+)</sup> (allow just charge) (1)

(If answer refers to m/z C.E. = 0)

Attract water molecules more strongly (1)

(d)  $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$  (1)

(allow incorrect or omitted use of [] for concentration of Al ions)

(Where A<sup>-</sup> is  $[\text{Al}(\text{H}_2\text{O})_5(\text{OH})]^{2+}$  and HA is  $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$ )

$K_a = \frac{[\text{H}^+]^2}{[\text{HA}]}$  when  $[\text{H}^+] = [\text{A}^-]$  therefore  $[\text{H}^+] = \sqrt{K_a[\text{HA}]}$  (1)

(this mark also scores the first mark)



$[\text{H}^+] = \sqrt{1.26 \times 10^{-5} \times 2.0} = 5.02 \times 10^{-3} \text{ (mol dm}^{-3}\text{)}$  (1)

pH = 2.30 (1)

(pH must be quoted to 2 d.p.) (QWC mark)

(Note pH = 2.30 scores 4)

(Note pH mark can be given consequentially on a wrong value for  $[\text{H}^+]$ )



(Note other equations possible)

pH = -1 to 1 (1)

**Total 13 marks**

**Question 9**

(a) Heat required = mass  $\times$  sp ht capacity  $\times$  rise in temp =  $1000 \times 4.18 \times 80 = 334400 \text{ J}$  (1)

(allow 334000 to 335000 J or 334 to 335 kJ)

Number of moles of methanol required to provide this =  $334400 / (715 \times 1000)$  (1)

(method mark for heat/(enthalpy of combustion) but both values must be in the same units)

= 0.4677 mol (1)

(allow 0.46 to 0.47)

But efficiency is only 0.5 therefore moles required =  $0.4677 \times 2 = 0.9354$  (1)

(note 0.935 scores 4)

(note this mark of 1 is for the factor of 2 and can be scored anywhere in the answer even if the rest of the calculation is wrong)

Mass = moles  $\times M_r = 0.935 \times 32 = 29.9 \text{ g}$  (1)

(allow 29 to 30.1 g allow answers to 2 sig figs)

(note correct answer scores 5)

(note answer of 14.5 to 15.1 scores 4/5)

- (b)  $K_c = [CH_3OH]/[H_2]^2[CO]$  (1)
- Moles at equilibrium of  $H_2 = 0.4$  (1)
- $CO = 0.2$  (1)
- Concentration = moles/vol = moles/1.5 (1)
- (can score this from next mark also)
- $K_c = (0.8/1.5) / (0.4/1.5)^2 \times (0.2/1.5)$  (1)
- 56.25 (1)

(allow 55.5 to 56.5)

(note correct answer scores 6)

(note an answer of 25 (not divided by vol to get concentration) scores 3/6)

$\text{mol}^{-2}\text{dm}^6$  (1)

(note mark units independently)

(Note if moles of  $H_2$  wrong and moles CO wrong, max mark is 3 for

$K_c$  expression, moles/ vol expression for concentration and units)

- (c) **Methyl ethanoate:** 2 peaks (1)
- Each is a singlet (1)
- Ethyl methanoate:** 3 peaks (1)
- Singlet, triplet, quartet all three scores 2 marks, ( or two out of three 1 mark) (1)

(Note must give number of peaks to score next mark(s)) (QWC)

(But if number of peaks can be unambiguously implied from splitting answer can score 1/2 for number of peaks (2 peaks then 3 peaks))

**Total 17 marks**