


UNIT 9

METALS AND THEIR COMPOUNDS

Student Version



Source: www.nde-ed.org

Contents

- Classification – Metals, Non-metals and Metalloids
- Physical Properties of Metals
- Chemical Properties of Metals
- Extraction of Metals
- Uses of Metals and Metalloids
- Properties and Reactions of Metal Compounds
- Qualitative Analysis Part 4

Key words: metal, non-metal, metalloid, malleability, ductility, lustre, sonority, alloy, bronze, brass, steel, solder, malleable, ductile, sonorous, lustrous, ferromagnetic, s-block metal, p-block metal, d-block metal, transition metal, rusting, reactivity series, metal displacement reaction, blast furnace, basic oxygen process, complex ion, ligand

Units which must be completed before this unit can be attempted:

- Unit 1 – Atoms and the Periodic Table
- Unit 2 – Particles, Bonding and Structure
- Unit 3 – Amount of Substance and Measurement
- Unit 4 – Introduction to Physical Chemistry
- Unit 5 – Acids, Bases and Salts
- Unit 6 – Redox Reactions
- Unit 8 – Solubility and Precipitation Reactions

Estimated Teaching Time: 10 hours

UNIT 9 – METALS AND THEIR COMPOUNDS

UNIT 9 SUMMARY AND SYLLABUS REFERENCE

Lesson	Title and Syllabus Reference
1	<p>Classification of Metals CA3ci metallic Bonding - factors influencing its formation (Factors should include: atomic radius, ionization potential, and number of valence electrons, type of specific packing not required); CA62bii alloys; ISA4.1 particulate nature of matter (elements - metals and non-metals (1st to 20th elements in the periodic table)); ISA10.1 classification of materials (classification of materials into metals, semi-metals (metalloids), and non-metals); CC8dii copper – properties; ISA10.2 uses of metals and semi-metals</p>
2	<p>Physical Properties of Metals CA2c elements of the first transition series (physical properties of the elements - physical properties should include: physical states, metallic properties and magnetic properties); CA3cii properties of metals (typical properties including conductivity - malleability, ductility demonstrated using metals like Mg, Zn, Sn, Fe); CC8a properties of sodium; CC8b properties of calcium; CC8dii copper – properties; ISA10.2 uses of metals and semi-metals (physical properties of metals and semi-metals under conductivity, luster, malleability, ductility, sonority, density, melting point and tensile strength)</p>
3	<p>Chemical Properties of Metals (s and p block) CA7a definitions of acids and bases (effects of acids and bases on metals); CA7b chemical properties of acids and bases (balanced chemical equations of all reactions); CC8a properties of sodium; CC8b properties of calcium; CC8c reactivity of aluminium with air, water and acids; ISA7.2 chemical properties of acids and bases (properties of acids and bases)</p>
4	<p>Chemical Properties of Metals (d block) CA2c elements of the first transition series (chemical reactivities of the elements, other properties should include variable oxidation states, catalytic abilities); CA7a definitions of acids and bases (effects of acids and bases on metals); CA7b chemical properties of acids and bases (balanced chemical equations of all reactions); CC8c reactivity of iron with air, water and acids; CC8dii copper – properties; ISA7.2 chemical properties of acids and bases (properties of acids and bases)</p>
5	<p>Reactivity Series of Metals CA2c elements of the first transition series (chemical reactivities of the elements); CA4aii chemical equations (displacement of metal ions); CC8a properties of sodium; CC8b properties of calcium; CC8c reactivities of aluminium and iron with air, water and acids; CC8dii copper – properties; ISA7.2 chemical properties of acids and bases (properties of acids and bases)</p>
6	<p>Extraction of Metals CA12bi extraction of metals - Al, Fe, Au and Sn (raw materials, processing, main products, byproducts, recycling)</p>
7	<p>Uses of Metals; Compounds of Sodium and Calcium CA12bi extraction of metals (uses of the metals); CA12bii alloys (common alloys of Cu, Al, Pb, and Fe and their uses); CC8a properties and uses of compounds of sodium (NaCl, NaOH, NaNO₃ and Na₂SO₄); CC8b properties and uses of compounds of calcium (CaO, CaSO₄, CaCl₂, Ca(OH)₂); CC8diii copper – uses; ISA10.2 uses of metals and semi-metals (uses of the following elements: Al, Cu, Fe, Au, application of semi-metals); ISA10.3 alloys (examples of alloys and their constituent elements (steel, bronze, brass), uses of alloys, advantages of alloys in manufacture of certain household items)</p>
8	<p>Compounds of D-block Metals CA2c elements of the first transition series (electronic configuration, chemical reactivities of the compounds, other properties should include formation of coloured compounds, complex ion formation); CA13ci characteristic tests of the following cations with dilute NH₃(aq): Ca²⁺, Pb²⁺, Cu²⁺, Fe²⁺, Fe³⁺, Al³⁺ and Zn²⁺; CC7b water and solution (test for water will be required); CC8diii copper – uses of compounds (the compounds must be limited to CuSO₄, CuO and CuCl₂)</p>
9	<p>Qualitative Analysis – Complex Formation Reactions CA2c elements of the first transition series (chemical reactivities of the compounds, other properties should include complex ion formation); CA13ci characteristic tests of the following cations with dilute NH₃(aq) and NaOH: Ca²⁺, Pb²⁺, Cu²⁺, Fe²⁺, Fe³⁺, Al³⁺ and Zn²⁺, confirmatory tests for these cations</p>
10	<p>Unit 9 Revision and Summary</p>

UNIT 9 – METALS AND THEIR COMPOUNDS

The Periodic Table of the Elements

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
	6.9 Li lithium 3	9.0 Be beryllium 4	1.0 H hydrogen 1	10.8 B boron 5	12.0 C carbon 6	14.0 N nitrogen 7	16.0 O oxygen 8	19.0 F fluorine 9	20.2 Ne neon 10										4.0 He helium 2
	23.0 Na sodium 11	24.3 Mg magnesium 12		27.0 Al aluminium 13	28.1 Si silicon 14	31.0 P phosphorus 15	32.1 S sulfur 16	35.5 Cl chlorine 17	39.9 Ar argon 18										
	39.1 K potassium 19	40.1 Ca calcium 20		47.9 Ti titanium 22	50.9 V vanadium 23	52.0 Cr chromium 24	54.9 Mn manganese 25	55.8 Fe iron 26	58.7 Ni nickel 28	63.5 Cu copper 29	65.4 Zn zinc 30	69.7 Ga gallium 31	72.6 Ge germanium 32	74.9 As arsenic 33	79.0 Se selenium 34	79.9 Br bromine 35	83.8 Kr krypton 36		
	85.5 Rb rubidium 37	87.6 Sr strontium 38		91.2 Zr zirconium 40	92.9 Nb niobium 41	96.0 Mo molybdenum 42	[98] Tc technetium 43	101.1 Ru ruthenium 44	106.4 Pd palladium 46	107.9 Ag silver 47	112.4 Cd cadmium 48	114.8 In indium 49	118.7 Sn tin 50	121.8 Sb antimony 51	127.6 Te tellurium 52	126.9 I iodine 53	131.3 Xe xenon 54		
	132.9 Cs caesium 55	137.3 Ba barium 56		178.5 Hf hafnium 72	180.9 Ta tantalum 73	183.8 W tungsten 74	186.2 Re rhenium 75	190.2 Os osmium 76	195.1 Pt platinum 78	197.0 Au gold 79	200.6 Hg mercury 80	204.4 Tl thallium 81	207.2 Pb lead 82	209.0 Bi bismuth 83	[209] Po polonium 84	[210] At astatine 85	[222] Rn radon 86		
	[223] Fr francium 87	[226] Ra radium 88		[267] Rf rutherfordium 104	[268] Db dubnium 105	[271] Sg seaborgium 106	[272] Bh bohrium 107	[270] Hs hassium 108	[281] Ds darmstadtium 110	[280] Rg roentgenium 111	Elements with atomic numbers 112-116 have been reported but not fully authenticated								
				140.1 Ce cerium 58	140.9 Pr praseodymium 59	144.2 Nd neodymium 60	[145] Pm promethium 61	150.4 Sm samarium 62	152.0 Eu europium 63	157.3 Gd gadolinium 64	158.9 Tb terbium 65	162.5 Dy dysprosium 66	164.9 Ho holmium 67	167.3 Er erbium 68	168.9 Tm thulium 69	173.1 Yb ytterbium 70	175.0 Lu lutetium 71		
				232.0 Th thorium 90	231.0 Pa protactinium 91	238.0 U uranium 92	[237] Np neptunium 93	[244] Pu plutonium 94	[243] Am americium 95	[247] Cm curium 96	[247] Bk berkelium 97	[251] Cf californium 98	[252] Es einsteinium 99	[257] Fm fermium 100	[258] Md mendelevium 101	[259] No nobelium 102	[262] Lr lawrencium 103		

* 58 – 71 Lanthanides

† 90 – 103 Actinides

Lesson 1 – What are metals?

a) Classification – Metals, Non-metals and Metalloids

**Summary Activity 1.1: What can you remember about metals?**

- What is a metal?
- Describe the structure and bonding in a typical metal
- What are the properties of a typical metal?
- What is a non-metal?
- Describe the structure and bonding in a typical non-metal
- Why are some elements metals and others non-metals?
- What is an alloy? How many alloys can you name?

(i) Metals

- **Metals** are elements which contain metallic bonding
- In metallic bonding, the atoms lose some or all of their outer shell (valence) electrons to form a lattice of cations and a sea of delocalised electrons; the delocalised electrons and the cations are attracted to each other and this holds the structure together
- Metallic bonding is preferred when all of the bonding atoms are electropositive, and hence do not hold on tightly to their valence electrons; it is therefore most commonly found in the elements of atoms on the left-hand side, and at the bottom, of the Periodic Table (bonding in elements becomes less metallic and more covalent from left to right across a Period, and more metallic and less covalent from top to bottom in a Group)
- In some cases, two or more different atoms can form a metallic bond; usually if both atoms are electropositive (ie come from metals; eg Cu and Zn); electropositive atoms can also form metallic bonds with atoms which are not electropositive (ie come from non-metals), provided that the electropositive atom is more abundant (eg Fe and C); mixtures which contain metallic bonds are known as **alloys**; alloys are not compounds because they do not have a fixed composition

(ii) Non-metals

- **Non-metals** are elements which do not contain metallic bonds; the vast majority of non-metals contain covalent bonding
- In elements, covalent bonding is preferred to metallic bonding if the atoms are not electropositive enough to give up their valence electrons; it is therefore most commonly found in the elements of atoms on the right-hand side, and at the top of the Periodic Table (bonding in elements becomes less metallic and more covalent from left to right across a Period, and more metallic and less covalent from top to bottom in a Group); elements containing covalent bonds are called **non-metals**

UNIT 9 – METALS AND THEIR COMPOUNDS

(iii) Metalloids

- **Metalloids** or **semi-metals** are substances which contain bonding intermediate in character between metallic and covalent; metalloids can be elements, compounds or mixtures, but the term is most commonly used to describe elements; often these substances display metallic properties at higher temperatures and non-metallic properties at lower temperatures; they are found in between the metals and the non-metals in the Periodic Table
- The following diagram shows the location of the metals, non-metals and metalloids in the Periodic Table; note the trend from metal to metalloid to non-metal across a Period from right to left and up a Group from bottom to top: (white = metal, grey = metalloid; black = non-metal)

I	II											III	IV	V	VI	VII	0			
		H																He		
Li	Be											B	C	N	O	F	Ne			
Na	Mg											Al	Si	P	S	Cl	Ar			
K	Ca	Sc				Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	Y				Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ba	La	Ce - Lu		Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn	
Fr	Ra	Ac	Th - Lw																	

(iv) Further classification of metals

- Metals can be further classified into s-block, p-block, d-block or f-block metals based on their electronic configuration and consequent position in the Periodic Table:
- Metals made from atoms containing valence electrons in s-orbitals only are called s-block metals (eg Na, Ca)
- Metals made from atoms containing valence electrons in s and d-orbitals only are called d-block metals (eg Fe, Cu, Au)
- Metals made from atoms containing valence electrons in s and p-orbitals only are called p-block metals (eg Al)



Test your knowledge 1.2: Classifying metals, non-metals and metalloids

- Name two s-block metals
- Name two p-block metals
- Name two d-block metals
- Name two metalloids
- Name two non-metals
- Why does metallic character decrease across a Period?
- Why does metallic character increase down a Group?

Lesson 2 – What are the physical properties of metals?

b) Physical Properties of Metals

- Most physical properties of metals are caused by the metallic bonding; in most cases, variations in the physical properties of the metals are caused by variations in the strength of metallic bonding:

(i) Strength of Metallic Bonding

- The strength of metallic bonding depends on:
 - The charge on the cations: the higher the charge on the cations, the stronger the metallic bonding
Eg Mg^{2+} is smaller than Na^+ so the metallic bonding is stronger in Mg than in Na
 - The size of the cations: the smaller the size of the cations, the stronger the metallic bonding
Eg Na^+ is smaller than K^+ so the metallic bonding is stronger in Na than in K
 - The number of delocalised electrons per atom: the greater the number of delocalised electrons, the stronger the metallic bonding; the number of delocalised electrons per atom is equal to the charge on the cation
Eg Mg gives up two electrons, Na gives up only one, so the metallic bonding is stronger in Mg than in Na
 - The block of the Periodic Table; the bonding in d-block metals is much stronger than the bonding in s and p-block metals (the reasons for this will not be discussed in this course)
- Generally, the further to the right, and the higher up, the element is found in the Periodic Table, the stronger the metallic bonding

(ii) Physical Properties of Metals

- Some physical properties are common to all metals:
 - **Electrical conductivity:** all metals contain delocalised electrons, which can move when a potential difference is applied; all metals are therefore good conductors of electricity
 - **Thermal conductivity:** the delocalised electrons carry heat energy with them, which can therefore move rapidly through the structure as the electrons move; all metals are therefore good conductors of heat
 - **Malleability:** in metals, the attraction between the delocalised electrons and the cations does not depend on the position of the cations; it is therefore possible to move the cations around without having a significant impact on the bonding; this means that metals can be easily bent into different shapes; all metals are therefore malleable
 - **Ductility:** because metals can be bent into different shapes, they can also be turned into wires; this property is known as ductility; all metals are therefore ductile, especially those with stronger metallic bonding
 - **Lustre:** the delocalised electrons in metals reflect light back towards its source; this causes metals to have a characteristic shiny appearance known as lustre
 - **Sonority:** the delocalised electrons in metals can also transport sound energy efficiently through the solid, which means that the sound wave can travel through the metal with little loss of energy; the ability of a solid to make a deep, ringing sound when hit is known as sonority; most metals are sonorous (although high density metals are less sonorous)

UNIT 9 – METALS AND THEIR COMPOUNDS

- Some other physical properties of metals depend on the strength of the metallic bonding, which varies significantly from metal to metal:
 - **Melting point:** the melting point of metals depends on the strength of the metallic bonding; it is generally high as metallic bonding is generally strong, so a lot of energy is needed to break up the metallic lattice; however the strength of the metallic bonding varies widely between metals and so does the melting point. Group I metals have weak metallic bonding and a low melting point; d-block metals have strong metallic bonding and a high melting point
 - **Tensile strength:** because metals are malleable, they can absorb large quantities of energy without break and so tend to have a high tensile strength; however the tensile strength depends on the strength of the metallic bonding, which varies widely between metals; Group I metals have weak metallic bonding and a tensile strength; d-block metals have strong metallic bonding and a high tensile strength

- Some properties of metals do not result from either the delocalised electrons or the strength of the metallic bonding:
 - **Magnetic Properties:** some metals have their delocalised electrons arranged in such a way that they can align themselves to a magnetic field and attract a magnet; the ability to do this is known as **ferromagnetism**; only Fe, Co and Ni are ferromagnetic; ferromagnetism is caused by the presence of unpaired electrons in the atoms: Fe, Co and Ni have electronic configurations d^6 , d^7 and d^8 respectively, which means that they have four, three and two unpaired electrons respectively; however not all atoms with unpaired electrons are ferromagnetic and the full causes of ferromagnetism will not be discussed in this course)
 - **Density:** density is mass per unit volume; the density of a metal therefore depends on the atomic mass (the higher the atomic mass the higher the density) and the atomic size (the larger the atomic size the lower the density); the density also depends to an extent on the type of crystal structure adopted by the metal, as some crystal structures are more compact than others; atomic mass increases across a Period and atomic size decrease across a period; the density of metals therefore increases across a Period; atomic mass increases down a group but so does atomic size; generally, the increase in atomic mass down a group is more significant so density tends to increase down a group; the metallic elements with the highest density are therefore to be found at the bottom right of the Periodic Table

- **Summary of properties of metals:**

Property	s-block and p-block	d-block
Electrical and Thermal Conductivity	High	high
Melting Point	low-medium	high
Tensile strength	low-medium	high
Malleability	High	high
Ductility	Medium	high
Lustre	High	high
Sonority	High	High
Magnetism	not-magnetic	Fe, Co, Ni magnetic
Density	low-medium	medium-high



Test your knowledge 2.1: Describing physical properties of metals

- Why do all metals conduct electricity?
- Why are all metals malleable?
- Why does magnesium have a higher melting point than sodium?
- Why does potassium have a lower melting point than potassium?
- Why is aluminium stronger than magnesium?
- Why is iron denser than aluminium?
- Why is iron more magnetic than copper?

Lesson 3 – How do s and p-block metals react with air, water and acids?



Summary Activity 3.1: What can you remember about redox reactions of metals?

- Give one example of a reaction between a metal and oxygen
- Give one example of a reaction between a metal and an acid
- Give one example of a metal displacement reaction
- In each reaction, what is oxidised and what is reduced?

c) Chemical Properties of Metals

(i) s-block metals (Na and Ca)

- s-block metals are electropositive and do not hold on to their electrons strongly; the valence electrons are easily lost during chemical reactions and the metal is oxidised; most s-block metals react rapidly with air, water and acids as a result
- s-block metals only form one stable cation, corresponding to the loss of all valence electrons; Group 1 metals form +1 ions only (eg Na⁺) and Group 2 metals form +2 ions only (eg Ca²⁺)
- Chemical reactivity increases as electropositivity increases; reactivity therefore decreases down a Group and Group 1 metals are more reactive than Group 2 metals
- With air, the elements will react to form oxides (containing O²⁻) (eg CaO)
 - although barium and all Group 1 metals except lithium can also form peroxides (containing O₂²⁻) (eg Na₂O₂)
 - these reactions are violent and exothermic once the metal is ignited
 - the more reactive s-block metals can ignite spontaneously, most s-block metals tarnish rapidly on exposure to air (they form an oxide layer at the surface of the metal which makes it less shiny)
 - $4\text{Na(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{Na}_2\text{O(s)}$; $2\text{Na(s)} + \text{O}_2\text{(g)} \rightarrow \text{Na}_2\text{O}_2\text{(s)}$; $2\text{Ca(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{CaO(s)}$
 - these reactions are redox reactions; the metal is oxidised and the oxygen is reduced; most s-block metals are stored under oil to prevent this reaction from taking place

UNIT 9 – METALS AND THEIR COMPOUNDS

- With water, the elements will react to form hydroxides (containing OH⁻) and hydrogen (H₂)
 - all s-block elements except Be and Mg react rapidly with water
 - with the most reactive metals the reaction is very explosive; this is another reason why the s-block metals are stored under oil; the metal fizzes rapidly as hydrogen is given off
$$2\text{Na(s)} + 2\text{H}_2\text{O(l)} \rightarrow 2\text{NaOH(aq)} + \text{H}_2\text{(g)}; \text{Ca(s)} + 2\text{H}_2\text{O(l)} \rightarrow \text{Ca(OH)}_2\text{(aq)} + \text{H}_2\text{(g)}$$
 - these reactions are redox reactions; the metal is oxidised and the hydrogen in the water is reduced
 - watch these reactions on: www.youtube.com/watch?v=fAFrzlyrqCU
 - with calcium, the solution quickly becomes cloudy; Ca(OH)₂ is only sparingly soluble in water and so the solution becomes saturated quickly; any more Ca(OH)₂ formed cannot dissolve and will form as a solid
- With acids, the reaction is even more vigorous and is dangerously explosive, especially with the Group 1 metals; the metal is oxidised and the H⁺ ion in the acid is reduced (to hydrogen gas); a salt is also produced (see Topic 6 – Redox Reactions)
$$2\text{Na(s)} + 2\text{HCl(aq)} \rightarrow 2\text{NaCl(aq)} + \text{H}_2\text{(g)}; \text{Ca(s)} + 2\text{HCl(aq)} \rightarrow \text{CaCl}_2\text{(aq)} + \text{H}_2\text{(g)}$$

(ii) p-block metals (Al, Sn and Pb)

- p-block metals are much less reactive than s-block metals; they are less electropositive and hold onto their electrons more strongly than s-block elements; of the common p-block metals, Al is the most reactive, followed by Sn and then lead
- aluminium only forms Al³⁺ ions, corresponding to the loss of all valence electrons; tin can lose all of its electrons to form Sn⁴⁺; tin can also lose its p-electrons only (not its s-electrons) to form Sn²⁺; lead generally loses its p-electrons only to form Pb²⁺
- Although Al is considered a reactive metal, it shows almost no reaction with air or water; the reaction is much slower than expected because a small quantity of aluminium at the surface can react to form Al₂O₃; this oxide forms a very stable layer on the surface of the metal which prevents any further oxygen or water from reacting with the aluminium
- Aluminium powder can react with air if heated, and aluminium can react with steam if heated strongly:
$$4\text{Al(s)} + 3\text{O}_2\text{(g)} \rightarrow 2\text{Al}_2\text{O}_3\text{(s)}$$
$$4\text{Al(s)} + 6\text{H}_2\text{O(g)} \rightarrow 2\text{Al}_2\text{O}_3\text{(s)} + 3\text{H}_2\text{(g)}$$
The aluminium is oxidised; with oxygen, the O is reduced; with water, the H is reduced
- Most p-block metals, including Al, will react slowly with acids to give a salt and hydrogen:
$$2\text{Al(s)} + 6\text{HCl(aq)} \rightarrow 2\text{AlCl}_3\text{(aq)} + 3\text{H}_2\text{(g)}$$
The Al is oxidised and the H⁺ in the acid is reduced (see Topic 6 – Redox Reactions)

UNIT 9 – METALS AND THEIR COMPOUNDS



Test your knowledge 3.2: Describing chemical properties of s and p-block metals

- Write equations for the reactions of potassium and calcium with water. Explain what you would observe, state the type of reaction taking place and explain why potassium reacts with water more quickly than calcium.
- Write equations for the reactions of sodium and calcium with oxygen. What is the difference between the two products?
- Why are the alkali metals stored under oil?
- Write equations for the reactions of magnesium and aluminium with hydrochloric acid. Explain the type of reaction taking place and explain why magnesium reacts with acids more quickly than aluminium
- Explain why aluminium shows little reaction with air or water despite being considered a reactive metal.

Lesson 4 - How do d-block metals react with air, water and acids?



Summary Activity 4.1: What can you remember about d-block metals?

- Why does iron rust?
- How can rusting be prevented?
- How is copper purified?
- What are the electronic configurations of Cu and Fe?
- What are the electronic configurations of Cu^{2+} , Fe^{2+} and Fe^{3+} ?

(iii) d-block metals (Fe, Cu and Au)

- d-block metals are generally less reactive than p-block metals, although their electropositivity and reactivity does vary considerably; of the common d-block metals, Zn is the most reactive, followed by Fe; Cu is unreactive; Ag and Au are very unreactive
- d-block metals always lose their valence s-electrons; most d-block metals have two s-electrons so form 2+ ions (eg Fe^{2+}) although atoms in columns 4 and 9 of the d-block only have one s-electron (eg Cu has the electronic configuration $[\text{Ar}]3d^{10}4s^1$) so can form +1 ions (eg Cu^+ , Au^+); the d-electrons in the outer shell, however, can behave as valence electrons in certain circumstances; this means that many atoms in the d-block can form several different ions, depending on how many d-electrons they lose:
 - Cu has the electronic configuration $[\text{Ar}]3d^{10}4s^1$; it loses its valence s-electron to form Cu^+ ; ($[\text{Ar}]3d^{10}$); it usually also loses one valence d-electron to form Cu^{2+} ($[\text{Ar}]3d^9$)
 - Fe has the electronic configuration $[\text{Ar}]3d^64s^2$; it loses its valence s-electrons to form Fe^{2+} ($[\text{Ar}]3d^6$); it can also lose one valence d-electron to form Fe^{3+} ($[\text{Ar}]3d^5$)

Transition metals can therefore have **several stable oxidation states**; it is common for d-block metals to form stable ions in which the d-orbitals are partially filled (ie ions which have 1 – 9 electrons in their d-orbitals); a metal which forms at least one stable ion with a partially filled d-orbital is called a **transition metal**; Cu, Fe and Au are all transition metals

UNIT 9 – METALS AND THEIR COMPOUNDS

- The ability of the d-electrons in transition metals to be lost and gained again results in transition metals (and their compounds) being good **catalysts**:
 - Fe is used as a catalyst in the Haber Process (the manufacture of ammonia: $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$)
 - Ni is used as a catalyst in the hydrogenation of alkenes ($\text{C}_2\text{H}_4 + \text{H}_2 \rightleftharpoons \text{C}_2\text{H}_6$)
- Reaction with air:
 - Most d-block metals show almost no reaction with air or water at normal temperatures, either because they are unreactive or because they form a stable oxide layer
 - Iron can react with air, when heated, to form iron (II) oxide and iron (III) oxide; copper can also react with air when heated to form copper (II) oxide; $2\text{Fe(s)} + \text{O}_2 \rightarrow 2\text{FeO}$; $4\text{Fe(s)} + 3\text{O}_2 \rightarrow 2\text{Fe}_2\text{O}_3$; $2\text{Cu} + \text{O}_2 \rightarrow 2\text{CuO}$
 - Gold does not readily react with oxygen; this is why gold remains shiny for a long time
- Reaction with water:
 - Iron does not readily react with water in the absence of air but in the presence of air, iron can react with both oxygen and water to form rust: $4\text{Fe} + 6\text{H}_2\text{O} + 3\text{O}_2 \rightarrow 4\text{Fe(OH)}_3$; unlike most other oxides and hydroxides of d-block elements, rust does not provide protect the iron surface and so iron will continue to rust if not protected
 - Copper is less reactive than iron and also forms a protective oxide layer; it does not react with water at all at room temperature but if it exposed to hot water and air for a period of time it can form a green layer (copper carbonate) on its surface; this layer also provides protection to the copper
 - Gold does not readily react with water
- Reaction with acids:
 - Some d-block metals, including Fe and Zn will react slowly with acids to give a salt and hydrogen: $\text{Fe(s)} + 2\text{HCl(aq)} \rightarrow \text{FeCl}_2\text{(aq)} + \text{H}_2\text{(g)}$; the Fe is oxidised and the H^+ in the acid is reduced; this can be a useful way of preparing salts of d-block metals
 - Cu and Au do not react readily with dilute acids



Test your knowledge 4.2: Describing chemical properties of d-block metals

- Explain why d-block metals often have more than one stable oxidation state
- What is the difference between a d-block metal and a transition metal?
- Explain why transition metals are good catalysts and give an example of a reaction catalysed by a transition metal
- Write the equation for the rusting of iron
- Explain why copper is used in water pipes and why gold makes good jewellery
- Write an equation for the reaction between iron and sulphuric acid

UNIT 9 – METALS AND THEIR COMPOUNDS

Lesson 5 – How can we compare the reactivity of different metals?



Summary Activity 5.1: Which types of redox reaction involve metals?

- Write an equation to show how sodium reacts with water; indicate what is oxidised and what is reduced; why does zinc not react with water?
- Write an equation to show how zinc reacts with hydrochloric acid; indicate what is oxidised and what is reduced; why does copper not react with hydrochloric acid?
- Write an equation to show how zinc reacts with copper sulphate; indicate what is oxidised and what is reduced; why does copper not react with zinc sulphate?
- Write an equation to show how zinc oxide reacts with carbon; indicate what is oxidised and what is reduced; why does aluminium oxide not react with carbon?
- Write an equation to show how copper oxide reacts with hydrogen; indicate what is oxidised and what is reduced; why does zinc oxide not react with hydrogen?
- What is the general name given to all of these reactions?

(iv) The Reactivity Series of Metals

- The reactivity of different metals to oxygen, metals and water varies widely; it is possible to rank the reactivity of the different metals (and some non-metals); this is known as the reactivity series; the metals at the top are the most reactive:

Metal	Reactivity
Sodium	Reacts with cold water
calcium	Reacts with cold water
magnesium	Reacts with acids but not water
aluminium	Reacts with acids but not water
(carbon)	
zinc	Reacts with acids but not water
iron	Reacts with acids but not water
tin	Reacts with acids but not water
lead	Reacts with acids but not water
copper	No reaction with acids or water
gold	No reaction with acids or water

- The higher up the reactivity series a metal is, the more reactive it is and the more vigorously and quickly it will react with water, acids, oxygen and other non-metals; in addition, the more reactive the metal is, the harder it is to extract the metal from its compounds
- More reactive metals can displace less reactive metals from their compounds; for example, zinc will displace copper from copper sulphate solution: $\text{Zn} + \text{CuSO}_4 \rightarrow \text{ZnSO}_4 + \text{Cu}$; these reactions are called **metal displacement** reactions; they also work in the solid state; for example, aluminium will reduce iron oxide to iron: $2\text{Al} + \text{Fe}_2\text{O}_3 \rightarrow 2\text{Fe} + \text{Al}_2\text{O}_3$

UNIT 9 – METALS AND THEIR COMPOUNDS

- These reactions are all examples of redox reactions (see Topic 6 – Redox Reactions); the reactivity of the metals can be deduced directly from the standard electrode potentials, the most common of which are shown below:

Half-reaction	Conventional representation	E°/V
$Mg^{2+}(aq) + 2e^- \rightleftharpoons Mg(s)$	$Mg Mg^{2+}$	-2.37
$Al^{3+}(aq) + 3e^- \rightleftharpoons Al(s)$	$Al Al^{3+}$	-1.66
$Zn^{2+}(aq) + 2e^- \rightleftharpoons Zn(s)$	$Zn Zn^{2+}$	-0.76
$Fe^{2+}(aq) + 2e^- \rightleftharpoons Fe(s)$	$Fe Fe^{2+}$	-0.44
$Pb^{2+}(aq) + 2e^- \rightleftharpoons Pb(s)$	$Pb Pb^{2+}$	-0.13
$2H^+(aq) + 2e^- \rightleftharpoons H_2(g)$	$Pt H_2, 2H^+$	0.00 (by definition)
$Cu^{2+}(aq) + 2e^- \rightleftharpoons Cu(s)$	$Cu Cu^{2+}$	+0.34
$Ag^+(aq) + e^- \rightleftharpoons Ag(s)$	$Ag Ag^+$	+0.80

- the more negative the standard electrode potential, the more reactive the metal; the above table shows clearly that the metals can be ranked in decreasing order of reactivity as follows: magnesium, aluminium, zinc, iron, lead, (hydrogen), copper, gold (see Unit 6 – Redox Reactions)



Practical 5.2: Compare the reactivity of different metals

- Using a measuring cylinder, pour 5 cm^3 of 0.5 mol dm^{-3} $ZnSO_4$ into each of three different test tubes
- Use a thermometer to measure the temperature of each solution
- Add a small spatula measure of magnesium powder to one test tube
- Observe what happens and note any temperature change
- Repeat steps (c) to (d) but this time adding copper powder instead of magnesium powder
- Repeat steps (c) to (d) but this time adding iron powder instead of magnesium powder
- Repeat steps (a) to (f) but this time use 5 cm^3 of 0.5 mol dm^{-3} $CuSO_4$, and add zinc powder to the first test tube, magnesium powder to the second test-tube and iron powder to the third test tube
- Repeat steps (a) to (f) but this time use 5 cm^3 of 0.5 mol dm^{-3} $FeSO_4$, and add zinc powder to the first test tube, magnesium powder to the second test-tube and copper powder to the third test tube
- Repeat steps (a) to (f) but this time use 5 cm^3 of 0.5 mol dm^{-3} $MgSO_4$, and add zinc powder to the first test tube, copper powder to the second test-tube and iron powder to the third test tube
- Record your results in a table similar to the one below:

Metal powder	salt solution			
	$ZnSO_4$	$CuSO_4$	$FeSO_4$	$MgSO_4$
Zn				
Cu				
Fe				
Mg				

**In which mixtures does a reaction take place? Write an equation for each reaction.
Can you list the four metals copper, magnesium, zinc and iron in order of reactivity?**



Test your knowledge 5.3: Understanding metal displacement reactions

Predict whether a chemical reaction will take place when the following substances are mixed; write the symbol equation for any reaction occurring:

- (a) copper and magnesium sulphate
- (b) magnesium and copper sulphate
- (c) magnesium sulphate and zinc
- (d) zinc sulphate and copper
- (e) zinc and copper sulphate
- (f) aluminium oxide and carbon
- (g) iron (III) oxide and carbon
- (h) tin (IV) oxide and carbon

Lesson 6 - How are metals extracted from their ores?



Summary Activity 6.1: Electrolytic Processes

- Give the half-equations for the reactions occurring at each electrode when Al is extracted from Al_2O_3
- Give the half-equations for the reactions occurring at each electrode during the purification of copper

d) Extraction of Metals

(i) General Principles of Extraction

- A mineral containing a sufficient quantity of a metal for its extraction to be commercially viable is called an **ore**
- Because metals are so useful, the extraction of metals from their ores is very commercially important; there are a number of different processes by which metals can be extracted from their ores; the choice of extraction process depends on:
 - the position of the metal in the reactivity series
 - the cost of the process
 - the required purity of the metal
- Metals can be extracted by electrolysis:
 - this works for all ores in which the bonding is ionic
 - it produces a pure form of the metal
 - but electrolysis is very expensive and only used for the most reactive metals
 - **aluminium**, for example, is extracted by the electrolysis of purified bauxite (aluminium oxide)
- Metals can also be extracted by reacting them with another element which is more reactive:
 - carbon is often used for this purpose as the process is very cheap
 - **iron, tin** and **copper** are extracted by reducing their ores (eg Fe_2O_3 and SnO_2) with carbon
 - this method cannot be used if the metal is above carbon in the reactivity series (like aluminium) or if the metal is not useful when contaminated with carbon (like titanium)
 - in such cases hydrogen or a reactive metal such as calcium should be used

UNIT 9 – METALS AND THEIR COMPOUNDS

- Metals which are present only in low concentrations require a more complex extraction process, and this is usually only commercially viable for very precious metals (like **gold**)

(ii) Extraction of aluminium

- Al_2O_3 is purified and then electrolysed (see Unit 6 – Redox Reactions):
 - the Al_2O_3 is dissolved in molten cryolite (Na_3AlF_6) as it has a lower melting point than Al_2O_3
 - the electrode reactions are: $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al}$ (at the cathode) and $2\text{O}^{2-} \rightarrow \text{O}_2 + 4\text{e}^-$ (at the anode)
 - graphite electrodes are used
 - the graphite anodes react with the oxygen produced and so they need to be regularly replaced
- This process uses a lot of energy and so is expensive; for this reason, recycling aluminium is very important

(iii) Extraction of iron

- Iron is extracted from its main ore haematite (Fe_2O_3) by reaction with carbon in a blast furnace; limestone is also added to remove the main impurity (SiO_2)
 - The carbon is added in the form of coke and it reacts as follows: $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$; $\text{C} + \text{CO}_2 \rightarrow 2\text{CO}$
 - the CO reduces the iron oxide to iron: $\text{Fe}_2\text{O}_3 + 3\text{CO} \rightarrow 2\text{Fe} + 3\text{CO}_2$
 - Some Fe_2O_3 may be reduced by C directly: $\text{Fe}_2\text{O}_3 + 3\text{C} \rightarrow 2\text{Fe} + 3\text{CO}$
 - limestone (CaCO_3) is also added; it decomposes on heating to give CaO: $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$
 - the CaO (which is basic) reacts with the main impurity SiO_2 (which is acidic): $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$
 - the CaSiO_3 is known as “slag”; it has a low density and can be separated from the reaction mixture; CaSiO_3 is used in road surfacing; it is laid underneath the tar
- The iron produced by this method contains a large quantity of carbon (up to 4%); some carbon is desirable (pure iron combined with a small quantity of iron makes the alloy steel) but 4% is too high; it is therefore necessary to blow oxygen through the molten iron in order to remove some of the C as CO and CO_2 ; other impurities such as sulphur can also be removed in this way; this is called the **basic oxygen process**
- the amount of carbon to be removed depends on the amount of carbon wanted in the steel; low-carbon steel is soft and used to make decorative items such as railings; medium-carbon steel is harder but still flexible and mainly used in construction; high-carbon steel is hard and brittle and used in tools
- iron is easy to recycle; it can be separated from other metals due to its magnetic properties

(iv) Extraction of tin

- Tin is reduced in a similar method to iron; its ore SnO_2 is heated with iron in a furnace: $\text{SnO}_2 + 2\text{C} \rightarrow \text{Sn} + 2\text{CO}$

(v) Extraction of gold

- Gold exists naturally as an element; its extraction consists of concentrating the gold by separating it from the surrounding rock
- Gold is unreactive but it can be converted into a soluble compound in the presence of oxygen and cyanide ions:

UNIT 9 – METALS AND THEIR COMPOUNDS

- This compound can be converted back into gold by reaction with carbon; the gold is formed on the surface of the carbon



Test your knowledge 6.2: Understanding the Extraction of Metals

- State three factors which determine how a metal is extracted from its ore
- Explain why the extraction of aluminium is expensive
- Explain why aluminium oxide is dissolved in molten cryolite
- Write the equations for the electrode reactions taking place during the electrolysis of aluminium oxide
- Explain why the anodes have to be regularly replaced
- Write two equations to show how carbon monoxide is produced in a blast furnace
- Write an equation for the reaction of tin (IV) oxide and carbon monoxide
- Write an equation for the reaction of iron (III) oxide and carbon monoxide
- Explain why limestone is added to the blast furnace
- Explain how excess carbon is removed from the iron produced in the blast furnace
- Explain why aluminium cannot be produced in the blast furnace
- State one useful by-product of the blast furnace and explain why it is useful
- Explain why iron is easier to recycle than aluminium or tin
- Explain why the extraction of gold is difficult, even though it occurs naturally
- Outline the stages in the extraction of gold (no equations required)

Lesson 7 – Why are metals and their compounds useful (part I)?

e) Uses of Metals and Metalloids

- The uses and applications of different metals depends on their physical and chemical properties:
 - **Sodium** is too reactive to be useful in elemental form; due to its reactivity, it can be used to produce titanium from titanium chloride; this is used in the extraction of titanium from its ore: $\text{TiCl}_4 + 4\text{Na} \rightarrow 4\text{NaCl} + \text{Ti}$
 - **Calcium** is also too reactive to be useful in elemental form; like sodium, its reactivity makes it useful in extracting other elements from their ores; calcium is used in the extraction of uranium from its ore
 - **Aluminium** is very useful in elemental form due to its strength, low density and resistance to corrosion (due to its protective oxide layer); it has a lower density than iron and doesn't corrode as easily; it is also non-toxic so can be used in food containers; aluminium is used in aeroplane parts, cans, foils and kitchen utensils; aluminium is often combined with other elements to make alloys which improve its properties (silicon is added to make it harder and stronger; magnesium is added to reduce its density)
 - **Tin** forms a protective oxide layer and is often used to coat other metals to prevent corrosion; tin combines with copper to make the alloy bronze; this is used to make statues and ornaments, as well as coins; tin combines with **lead** to make the alloy **solder**; this has a low melting point and is useful for welding electrical components together
 - **Iron** is very strong, especially when small amounts of carbon are added to make **steel**; steel is used to make steel girders for construction and is used in reinforced concrete; it is also used in bridges, pylons, bicycles, cars and pipes; because of its magnetic properties it is also used to make magnets
 - **Gold** is very unreactive and also soft; it is used in jewellery as it stays shiny for a long time; it is also stored as currency

UNIT 9 – METALS AND THEIR COMPOUNDS

- **Copper** is unreactive and very malleable so is used to make coins and pipes; it is a good conductor of electricity so it is used to make electrical wires; when mixed with zinc it makes the alloy **brass**; brass has antibacterial properties and it used in taps and door handles, especially in bathrooms; it is shiny and resists corrosion and is often used to coat ornaments; it is also highly sonorous and so used to make musical instruments (trumpets, trumbones, saxophones and horns are known as “brass” instruments); when copper is mixed with tin it makes the alloy **bronze**; bronze is very malleable and used in coins and sculptures; brass and bronze are both unreactive, like copper
- **Metalloids** (eg silicon and boron) are semi-conductors; they conduct poorly at low temperatures but better at high temperatures; for this reason they are very useful in making electronic components; they are also added to metals to make them harder (less metallic)



Test your knowledge 7.1: Describing uses of metals and metalloids

- (a) Give the name of the alloys formed by mixing the following elements:
(i) copper and tin; (ii) copper and zinc; (iii) tin and lead; (iv) iron and carbon
- (b) Complete the following table to show the at least one common use of the following metals and metalloids; in each case, identify the property which makes the material suitable for its use

Material	Use	Property
aluminium		
tin		
solder		
steel		
gold		
copper		
brass		
silicon		

f) Properties and Reactions of Metal Compounds

(i) Compounds of sodium

- All compounds of sodium are soluble in water; like most metals, it forms basic oxides and hydroxides and it can also form salts
- Sodium ions (Na^+) are colourless; they form colourless solutions and white solids, unless the anion in the compound is coloured; this is true of all s and p-block metal ions
- The most common salts of sodium are:
 - **sodium chloride (NaCl)** is a salt of a strong acid and a strong base; it is therefore neutral; it is commonly known as salt, common salt or table salt; it is added as flavouring to food and as a source of Na^+ ions; it is also placed on roads in cold weather to stop water freezing (all dissolved solutes lower the freezing point of water) as it is very cheap and non-toxic
 - **sodium nitrate (NaNO_3)** is a salt of a strong acid and a strong base; it is therefore neutral; it is commonly known as Chile saltpetre; it is an important source of nitrate ions which are important in fertilisers
 - **sodium sulphate (Na_2SO_4)** is a salt of a strong acid and a strong base; it is therefore neutral; it is mainly used in detergents
- **sodium hydroxide (NaOH)** is a strong base due to the OH^- ion; it is also known as caustic soda; it is mainly used to neutralise fatty acids to make soaps; it is also used in the manufacture of paper

UNIT 9 – METALS AND THEIR COMPOUNDS

(ii) Compounds of calcium

- Like most metals, calcium forms basic oxides and hydroxides and it can also form salts
- Calcium ions (Ca^{2+}) are colourless; they form colourless solutions and white solids, unless the anion in the compound is coloured; this is true of all s and p-block metal ions
- The most common salts of calcium are:
 - **Calcium chloride (CaCl_2)** is a salt of a strong acid and a strong base; it is therefore neutral; it is highly soluble in water even at low temperatures and is very effective at reducing the freezing point of water, hence preventing freezing; it is a very useful de-icer; it is also hygroscopic and so is used as a drying agent and is also placed on dusty roads so the road holds water better and gets less dusty, which preserves the road; it is also a water softener because the presence of Ca^{2+} ions stops CO_3^{2-} and SO_4^{2-} ions dissolving in water
 - **Calcium sulphate (CaSO_4)** is a salt of a strong acid and a strong base; it is therefore neutral; it is insoluble in water but is most stable in its hydrated form: $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ – known as gypsum; the anhydrous form (CaSO_4) and another hydrated form $\text{CaSO}_4 \cdot 0.5\text{H}_2\text{O}$ are both hygroscopic, which makes them useful as drying agents; $\text{CaSO}_4 \cdot 0.5\text{H}_2\text{O}$ is known as plaster of Paris – it forms a mouldable paste with water which hardens into $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ when it dries; this is the most useful application of CaSO_4
- **Calcium oxide (CaO)** is a base due to the O^{2-} ion; the O^{2-} ion is a strong base but CaO is only slightly soluble in water so is considered a weak base; CaO absorbs water readily to form $\text{Ca}(\text{OH})_2$: $\text{CaO} + \text{H}_2\text{O} \rightarrow \text{Ca}(\text{OH})_2$; it is therefore hygroscopic and a good drying agent; CaO is used to remove impurities in the manufacture of steel; it reacts with acidic oxides left in the iron (mainly SiO_2) to produce salts: eg $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$
- **Calcium hydroxide ($\text{Ca}(\text{OH})_2$)** is a base due to the OH^- ion; it is only slightly soluble in water and a saturated solution of $\text{Ca}(\text{OH})_2$ in water is known as limewater and is used to test for carbon dioxide; it is used in water treatment - it is added to water to reduce its acidity and the precipitate of $\text{Ca}(\text{OH})_2$ also acts as an effective water filter; $\text{Ca}(\text{OH})_2$ is used in the production of cement and in the production of paper; it can also be heated to convert it into CaO : $\text{Ca}(\text{OH})_2 \rightarrow \text{CaO} + \text{H}_2\text{O}$



Test your knowledge 7.2: Describing uses of the hydroxides, oxides and salts of sodium and calcium

- (a) Name the three most common sodium salts and state a use for each one
- (b) Name the two most common calcium salts and state a use for each one
- (c) State two uses for each of sodium hydroxide, calcium oxide and calcium hydroxide

Lesson 8 – Why are metals and their compounds useful (part II)?

(iii) Compounds of d-block metals (Cu, Zn and Fe)

- Most compounds of copper, zinc, iron and other d-block metals are bases or salts
- **Copper chloride (CuCl_2)** is a salt of a strong acid and a relatively weak base
 - it is therefore very slightly acidic due to the hydrolysis of the Cu^{2+} ion
 - it is a green solid and dissolves in water to form a blue-green solution
 - it is used as a catalyst in a variety of industrial processes



Test your knowledge 8.3: Describing properties and reactions of compounds of d-block metals

- (a) Give the formula of the two most common copper salts and give one use of each
- (b) Give the formula of the most common base of copper and state one use for it
- (c) Define the terms “complex ion” and “ligand”; give one example of a complex ion
- (d) Explain why complex ions of d-block metals are often coloured
- (e) Explain why zinc compounds are colourless
- (f) Explain why anhydrous copper sulphate is colourless
- (g) Write an equation for the reaction between Cu^{2+} and excess ammonia and state what you would see
- (h) Write an equation for the reaction between Zn^{2+} and excess ammonia and state what you would see
- (i) Describe a chemical test for water and write an equation for the reaction taking place

Lesson 9 - How can we use complex formation reactions in qualitative analysis?

g) Qualitative Analysis Part 4 – Complex Formation Reactions



Summary Activity 9.1: What can you remember about qualitative analysis?

- What is qualitative analysis?
- Which cations form precipitates with aqueous sodium hydroxide? What are the colours of these precipitates?
- Which of these precipitates dissolve in excess aqueous sodium hydroxide?

- Some simple qualitative tests for cations, anions and gases were covered in Unit 5 – Acids, Bases and Salts, Unit 6 – Redox Reactions and Unit 8 – Solubility and Precipitation
- Complex formation reactions are useful in qualitative analysis; Fe^{2+} , Fe^{3+} , Ca^{2+} , Al^{3+} , Pb^{2+} , Cu^{2+} and Zn^{2+} all give insoluble hydroxide precipitates when added to ammonia but only $\text{Cu}(\text{OH})_2$ and $\text{Zn}(\text{OH})_2$ dissolve in excess ammonia to give a complex ion

UNIT 9 – METALS AND THEIR COMPOUNDS



Practical 9.2: Use complex formation reactions to identify cations in solution

You are given four colourless solutions, labelled A, B, C and D; each solution contains one cation from the following: Ca^{2+} , Zn^{2+} , Pb^{2+} and Al^{3+}

- 1) Pour 2 cm³ of solution A into a test tube
- 2) Add a few drops of NaOH, then adding NaOH until the test tube is almost full; note your observations
- 3) Repeat steps (a) to (b) using solutions B – D
- 4) Pour 2 cm³ of solution A into a test tube
- 5) Add a few drops of NH₃, then adding NH₃ until the test tube is almost full; note your observations
- 6) Repeat steps (a) to (b) using solutions B – D

Use your observations to identify which cation is in which solution; present your answers in a table similar to the one below:

Solution	Observations				Cation Present
	Few drops NaOH	Excess NaOH	Few drops NH ₃	Excess NH ₃	
A					
B					
C					
D					

Which two cations cannot be distinguished by this combination of tests? How would you distinguish between solutions containing these cations?



Test your knowledge 9.3: Understanding qualitative analysis by complex formation reactions

- (a) Describe a chemical test to distinguish between solutions of aluminium sulphate and zinc sulphate

UNIT 9 – METALS AND THEIR COMPOUNDS

Lesson 10 – What have I learned about metals and their compounds?



10.1 END-OF-UNIT QUIZ UNIT 9 – METALS AND THEIR COMPOUNDS

1. Describe the bonding in metals and hence explain why metals are malleable and ductile.
2. State two other physical properties of metals and explain why metals have these properties.
3. Write equations for the following reactions and state the type of reaction occurring:
 - (a) Magnesium with dilute nitric acid
 - (b) Sodium with water
 - (c) Zinc with copper sulphate
4. Explain why zinc reacts with copper sulphate but copper does not react with zinc sulphate.
5. Describe how aluminium is extracted from its ore.
6. State why brass and solder are useful alloys and state which pure metals they are made from.
7. Explain why aqueous solutions of zinc salts are colourless but aqueous solutions of copper salts are coloured.
8. Explain the difference between a d-block metal and a transition metal.
9. Explain why transition metals can form variable oxidation states.
10. State what you would see when dilute ammonia solution is added gradually to a solution containing Zn^{2+} ions. Write equations for any reactions occurring.
11. Explain how you would distinguish between aqueous solutions of zinc nitrate and aluminium nitrate.
12. Describe a chemical test for water.